


Mitosis & Cytokinesis





- Mitosis: Dividing cell's DNA between 2 daughter nuclei


- 4 phases:

- prophase
- metaphase
- anaphase
- Telophase

- Cytokinesis: Dividing cell's cytoplasm



Overview of mitosis



green = key features



End of G₂ & Start of Mitosis

■ End G₂

- ◆ 2 Centrosomes have formed by replication of one centrosome
 - Each has 2 centrioles in animal cells
- ◆ Duplicated chromosomes cannot be seen yet individually under a microscope


■ Prophase (1st sub-phase of mitosis)

- ◆ **Chromatin condenses**
 - Starts turning into visible chromosomes
 - ◆ Each made of 2 sister chromatids
- ◆ **Nucleolus disappears**
- ◆ **Centrosomes move to opposite cell poles**
 - microtubules between them lengthen
 - ◆ Asters visible (radial arrays of short microtubules around centrosome)
- ◆ **Microtubules are forming the mitotic spindle**
 1. Coordinates movement of chromosomes
 2. Will elongate the cell later in mitosis



End of G₂ (of Interphase)


- ◆ Chromosomes were **replicated** during S phase of interphase
 - ◆ Each 1 chromosome is made up of 2 identical copies of double-helical DNA called **sister chromatids**



- ◆ Remember, earlier in G₂
 - ◆ DNA errors made during replication are corrected when possible
 - ◆ Organelles are duplicated, if needed
 - ◆ Proteins needed during M phase (mitosis & cytokinesis) are synthesized
 - ◆ Cell may grow slightly more, if necessary

Start of Mitosis (Prophase)

EARLY PROPHASE






Transitioning to Metaphase

green = key features

■ Prometaphase (*late prophase*)



- ◆ Nucleus fragments (breaks up)
- ◆ Chromosomes condense still more
- ◆ Two kinetochores appear on the side of each chromatid at the centromere of the duplicated chromosome
 - Chromosomes attach to spindle fibers at kinetochores
 - ◆ Structures of proteins located at the centromere of the chromosomes
- ◆ Spindle fibers are attaching to centromeres of chromosomes and become known as “kinetochore microtubules”



Transitioning to Metaphase

green = key features

- **Prometaphase continued** (*late prophase*)
 - ◆ Chromosomes begin moving toward a region equidistant between the two centrosomes, which are now located on opposite ends of the cell
 - ◆ “nonkinetochore microtubules” originating from one pole interact with those originating at the opposite pole of the mitotic spindle
 - Nonkinetochore microtubule interactions begin to cause some lengthening of the cell




Pro-metaphase (Late Prophase)


Centromeres

- Regions of DNA where the sister chromatids are most tightly connected via proteins
- Region where kinetochore proteins will attach to each sister chromatid

LATE PROPHASE
(PROMETAPHASE)





SPINDLE ANATOMY



green = key features

Metaphase (longest phase; 20min.)


- **Chromosomes align along middle of cell**
 - ◆ **On the metaphase plate**
 - meta = middle
 - ◆ Imaginary line half way between the spindle poles
 - ◆ **spindle fibers coordinate movement of chromosomes**
 - ◆ **Kinetochores microtubules from opposite poles are attached to the kinetochore of each sister chromatid in a chromosome**
 - ◆ **helps to ensure chromosomes separate properly**
 - so each new nucleus receives only 1 copy of **each** chromosome




Metaphase (longest phase; 20min.)

METAPHASE

metaphase plate is not a physical structure, just a term for the plane where the chromosomes line up.




← chromosomes line up at metaphase plate



(a) Diagram of two duplicated chromosomes arrayed at the metaphase plate



Metaphase:
Sister chromatids face opposite poles.



green = key features

Anaphase (shortest phase; few min.)


- **Sister chromatids separate**
 - **Cohesin** proteins, holding them together, are **cleaved**
 - ◆ **Each chromatid now considered an individual chromosome**
 - Each chromatid of a chromosome is pulled to opposite poles from the kinetochores at the centromeres
 - ◆ Pulled by motor proteins “walking” along microtubules (ATP)
 - Increased production of ATP by mitochondria
 - ◆ Kinetochores microtubules shorten
- **Poles move farther apart (cell lengthens)**
 - ◆ “non-kinetochore microtubules” lengthen push apart




Anaphase

- **Sister chromatids separate** (becoming independent **UNDUPLICATED** chromosomes)
 - ◆ Kinetochore microtubules pull sister chromatids to opposite poles by shortening
- **Poles of the cell are pushed farther apart** (cell lengthens)
 - ◆ Non-kinetochore microtubules elongate, interact via motor proteins, & push the centrosome/asters further apart, thereby, pushing the whole plasma membrane outward at both poles

ANAPHASE A CHROMOSOMES ARE PULLED POLEWARD





ANAPHASE B POLES MOVE APART




Separation of chromatids

- In anaphase, proteins holding together sister chromatids are inactivated
 - ◆ Chromatids separate to become individual chromosomes




Chromosome movement

- Kinetochores use motor proteins (dynein) that “walk” chromosome along attached microtubule
 - ◆ microtubule shortens by dismantling at kinetochore (chromosome) end




(a) Hypothesis



(A) ATP-driven motor protein drives both chromosome movement and microtubule disassembly

(B) microtubule disassembly drives chromosome movement




(b) Experiment


Alternative hypotheses for how microtubules shorten

Telophase - the end of mitosis

- Chromosomes arrive at opposite poles
 - ◆ 2 daughter nuclei form
 - Nuclei forms
 - Nucleoli form
 - ◆ chromosomes disperse (uncoil)
 - no longer visible under light microscope as they turn back into being chromatin
- Spindle fibers break down
- Cytokinesis begins
 - ◆ Cytoplasm divides
 - Involves Cleavage Furrow in animal cells & Cell Plate formation in plant cells.




TELOPHASE AND CYTOKINESIS



After Mitosis & Cytokinesis

- After mitosis & cytokinesis, the daughter cells are back in G₁ of Interphase
- During G₁...
 - ◆ the cells begin performing their normal metabolic functions
 - ◆ gene transcription & mRNA translation start up again
 - ◆ the cell grows into its "adult" size



Cytokinesis


In Animals:

- ◆ Involves cleavage.
- ◆ Constricting belt of actin microfilaments together with myosin proteins forms around the equator of cell


- cleavage furrow forms = shallow groove in the cell surface near location of past metaphase plate

- ◆ splits cell in two


- Looks like the tightening of a draw string
- Actin filaments interact with plasma membrane proteins pulling the plasma membrane inward.




Mitosis in whitefish blastula




Interphase




Prophase




Metaphase



Anaphase






Early Telophase



Late Telophase

Mitosis in animal cells



Cytokinesis in Plants


■ Plants

◆ cell plate forms late in telophase


- vesicles line up at equator (move along microtubules)
 - ◆ *derived from Golgi*
- vesicles fuse together to form the eventual two cell membranes of the daughter cells.
- This growing line of fused vesicles [cell plate] eventually fuse with the plasma membrane along the perimeter of the dividing parent cell
- Contents of vesicle make up middle lamella (pectin-rich)
 - ◆ This is a carbohydrate “sticky” substance that holds plant cells together.

◆ Cellulose [the new primary cell walls] will then be secreted by the two new daughter cells directly from the plasma membrane enzyme *cellulose synthase* [not through exocytosis by vesicles from the Golgi]


- This new cell wall forms under the middle lamella, between the middle lamella and each plasma membranes
 - ◆ new cell wall forming in between the daughter cells fuse with existing cell wall on the other three sides of each daughter cell



Vesicles with Pectin fusing to form cell plate



Mitosis in plant cell



Interphase




Prophase



Prometaphase



Metaphase



Anaphase



Telophase

Cytokinesis

MITOSIS

Prophase



Metaphase

Anaphase

Telophase


- **Cytokinesis, the division of the cytoplasm, overlaps with the final stages of mitosis.**
 - ◆ **It may start in either anaphase or telophase, and finishes shortly after telophase.**
 - In animal cells, cytokinesis is contractile. The cell is pinched in two with the help of filaments made of actin proteins interacting with plasma membrane proteins.
 - Plant cells have a stiff cell wall that cannot be pinched in two. A cell plate forms from vesicles aligning down the middle of the cell, splitting it into two daughter cells separated by a new wall.
 - ◆ **When cytokinesis finishes, two new cells have formed, each with a complete set of chromosomes identical to those of the mother cell.**
 - **The daughter cells may one day undergo mitosis themselves, repeating the cycle.**

CYTOKINESIS



Binary Fission

- Mitosis in eukaryotes likely evolved from binary fission in bacteria (**prokaryotes**)
- ◆ Binary fission means ‘division in half’
- Prokaryotes do **NOT** use mitosis
- ◆ In bacteria, most genes are carried on a **SINGLE** bacterial chromosome consisting of a **circular DNA** molecule.
 - This single circular chromosome is much shorter than the multiple linear chromosomes of eukaryotes and so is faster to duplicate
 - No membrane-bound organelles exist, including no nucleus that surrounds the bacterial DNA



Evolution of mitosis

- A possible progression of mechanisms intermediate between binary fission & mitosis seen in modern organisms

Prokaryotes (bacteria)


- ✓ No nucleus
- ✓ Single circular chromosome. After DNA is replicated, it is partitioned in the cell.
- ✓ After cell elongation, proteins assemble into a ring and facilitates separation and cell division.



prokaryotes (bacteria)

Single Celled Protists like Dinoflagellates


- ✓ Chromosome attached to nuclear envelope
- ✓ Microtubules pass through nucleus
- ✓ Nucleus divides similar to binary fission





Protists: dinoflagellates

Single Celled Protists, Diatoms, & Yeasts

- ✓ Nuclear envelope remains intact
- ✓ Microtubules in yeast form spindle inside nucleus (in diatoms it forms between two pairs of centrioles and passes through a tunnel)
- ✓ Microtubules separate chromosomes
- ✓ Nucleus splits into daughter nuclei




protists diatoms



eukaryotes yeast

Most Eukaryotes

- ✓ Spindle forms outside the nucleus
- ✓ Nuclear envelope breaks down during mitosis
- ✓ Microtubules separate the chromosomes
- ✓ Nuclear envelope reforms



eukaryotes animals

Any Questions??

